

Macro algae as substrate for biogas production

Møller, Henrik; Sarker, Shiplu; Gautam, Dhan Prasad; Bruhn, Anette; Fredenslund, Anders Michael

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Møller, H., Sarker, S., Gautam, D. P., Bruhn, A., & Fredenslund, A. M. (2012). *Macro algae as substrate for biogas production*. Abstract from Nordic Biogas Conference, København, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

MACRO ALGAE AS SUBSTRATE FOR BIOGASPRODUCTION

Henrik B. Møller¹, Shiplu Sarker¹, Chitra S. Raju¹, Dhan Prasad Gautam¹
Anette Bruhn², Anders M. Fredenslund³
*Aarhus University, ¹Department of Engineering,
²Department of Bioscience, ³University of Roskilde*



AU RESEARCH FROM LAB TO FULL SCALE



ALGAE SPECIES TESTED FOR BIOGAS

Brunalger (*Phaeophyceae*)



Fingertang (*Laminaria digitata*)



Sukkertang (*Saccharina L*)



Søsalat (*Ulva lactuca*)




Krølhårstang (*Chaetomorpha linum*)



Buletang (*Ascophyllum nodosum*)




WHY ALGAE FOR BIOGAS




HARVEST POTENTIAL


Biomass yield



BIOGAS EXPERIMENTS-BATCH




Methane potentials




| | Laminaria | Saccharina | Chaetomorpha Linum | aschophyllum | Ulva lactua | Pilayella littoralis and Ectocarpus Siliculosus |
|---------|-----------|------------|-----------------------|--------------|-------------|--|
| 90 days | 7 | 2 | 7 | 8 | 24 | 0 |
| 60 days | 34 | 12 | 18 | 54 | 50 | 1 |
| 30 daYS | 257 | 206 | 162 | 119 | 100 | 117 |

METHANE POTENTIALS



SEASONAL VARIATION OF BMP IN LAMINARIA




From Adams et al. 2011


CONTINUOUSLY EXPERIMENTS – CODIGESTION OF MANURE AND ALGAE

| | <i>Ulva</i> | | | <i>Laminaria</i> | | | Control |
|--------------------|-------------|-------|-------|------------------|-------|-------|---------|
| %VS (algae:manure) | 11:89 | 24:76 | 37:63 | 15:85 | 24:76 | 41:59 | 0:100 |
| HRT, days | 22 | | | 22 | | | 22 |
| Temp.(°C) | 50 | | | 50/35 | | | 50/35 |

RESULTS OF CONTINUOUSLY THERMOPHILIC EXPERIMENTS




Ulva lactuca





Laminaria

FERTILIZER VALUE – DIGESTATE USE FOR ORGANIC FARMING



FERTILIZER AND HEAVYMETALS



OFF SHORE HARVEST/CULTIVATION OR CULTIVATION IN BASINS

Line cultivation



Basin cultivation



Sea harvest




Beach 'harvest'



ECONOMICS

- › Production costs: **3 kr/kg DM**
- › Methane value: **1-2 kr/kg DM**
- › **How to make it profitable:**
- › Value of nitrogen for organic farming:
- › **0,5 kr/kg DM (15 kr/kg N)**
- › Payment for N removal from water environment: **>75 kr/kg N**
- › Extraction of products of high value and only using the by product for biogas
- › Value on the removal from beaches etc. **??**

EXAMPLE OF A COMING BIOGAS PLANT BASED ON ALGAE - SOLRØD



BIOGAS PLANT BASED ON ALGAE - SOLRØD



The process is very stable, Methane Production >260 L/kg VS or 1.61 l/l of reactor per day, Methane content : 59%

CONCLUSIONS

- › Algae is a very high yielding biomass that will not compete with food production. Ulva can have a drymatter yield that is more than 10 times higher than crops
- › Laminaria was found to be the macro algae with highest gaspotential 250-300 L CH₄/kg VS.
- › Co-digestion of algae with manure is advantageous but the ratio that can be used depends on algae species, Laminaria is superior
- › Digestate is a high value fertilizer with low heavy metal content
- › Cultivation of algae for biogas is not economical feasible today, but valuing environmental benefits, extraction of high value components or collection of cast seaweed at beaches might be ways to integrate algae in biogasplants as in Solrød



THANKS FOR YOUR
ATTENTION